arXiv:hep-ph/9501236v2 18 Jan 1995

Ochanomizu University

OCHA-PP-46
YCCP-9408
November 5, 1994

Majorana neutrino versus Dirac neutrino in $e^+e^- \rightarrow W^+W^-$ through radiative corrections

Y. Katsuki, M. Marui, R. Najima*,
J. Saito** and A. Sugamoto

Department of Physics, Faculty of Science, Ochanomizu University,
Bunkyo-ku Tokyo 112, Japan

* *Department of General Education, Yokohama College of Comerce,*
Kanagawa 203, Japan

** *Suzugamine Women's College,*
Hiroshima 733, Japan

Abstract

Radiative corrections to $e^+e^- \rightarrow W^+W^-$ from Majorana neutrinos are studied in the context of the see-saw mechanism. Focusing on the effects of the fourth generation neutrinos, we calculate W-pair form factors, the differential cross sections and the forward-backward asymmetries for the polarized electrons at one-loop level. The behaviour of the form factors at the threshold of Majorana particle pair productions is found to differ from that of Dirac particle pair productions. In the cross section for unpolarized electrons, the radiative corrections, depending on the mass parameters of the see-saw mechanism, are found to be $\sim 0.5\%$ at the energy range of the LEP200 and the next generation linear colliders.

Submitted to *Physics Letters* **B**

Precision experiments can probe new physics beyond the standard model, because heavy particles do not necessarily decouple in low-energy processes [1]. In the processes $\mathrm{e^+e^-} \to \mathrm{f\overline{f}}$ observed at the current colliders (LEP, SLC), the dominant effects of heavy particles appear only in the corrections to the self erergies of gauge bosons. These effects can be summarized into three parameters S, T and U [2] or ϵ_1, ϵ_2 and ϵ_3 [3], which have been estimated in various models. On the other hand , in the process $\mathrm{e^+e^-} \to \mathrm{W^+W^-}$ which is the most important process in the LEP200 experiments, the effects of heavy particles appear not only through the self energies of gauge bosons but also through the trilinear gauge vertices $\mathrm{AW^+W^-}$ and $\mathrm{ZW^+W^-}$. Radiative corrections to this process in the standard model with heavy Dirac fermions or heavy scalars have already been studied [4]. In this letter we investigate radiative corrections to $\mathrm{e^+e^-} \to \mathrm{W^+W^-}$ from heavy Majorana neutrinos in the context of the see-saw mechanism [5]. Focusing on the fourth generation neutrinos, we calculate W-pair form factors and the differential cross sections for the polarized electrons and W bosons at one-loop level.

We consider a model containing the fourth generation leptons having no mixing with other generations [6]. Recent measurements at LEP and SLC do not rule out the existence of the fourth neutrino but constrain its mass to be greater than 45 GeV. We assume the following mass term for the neutrinos

$$\mathcal{L}_{\mathrm{mass}} = -\frac{1}{2}\overline{(N_{\mathrm{L}}\ N_{\mathrm{R}}^{\mathrm{c}})}\begin{pmatrix} 0 & m_{\mathrm{D}} \\ m_{\mathrm{D}} & M_{\mathrm{R}} \end{pmatrix}\begin{pmatrix} N_{\mathrm{L}}^{\mathrm{c}} \\ N_{\mathrm{R}} \end{pmatrix} + \mathrm{h.c.},$$

where $N^{\mathrm{c}} = C\overline{N}^{\mathrm{T}}$ is the charge-conjugated state of N. We expect that the Dirac masses of neutrinos m_{D} generated by the vacuum expectation value of ordinary Higgs bosons are comparable to the masses of their charged lepton partners M_{E}. On the other hand, the right-handed Majorana mass M_{R} originates from "beyond the standard model". In order to get positive mass eigenvalues, we diagonalize the mass matrix and perform the chiral transformation. The result is given by

$$\mathcal{L}_{\mathrm{mass}} = -\frac{1}{2}\overline{(N_1\ N_2)}\begin{pmatrix} M_1 & 0 \\ 0 & M_2 \end{pmatrix}\begin{pmatrix} N_1 \\ N_2 \end{pmatrix},$$

where

$$\begin{aligned} \begin{pmatrix} N_1 \\ N_2 \end{pmatrix} &= \begin{pmatrix} i\gamma_5 c & -i\gamma_5 s \\ s & c \end{pmatrix}\begin{pmatrix} N_{\mathrm{L}} + N_{\mathrm{L}}^{\mathrm{c}} \\ N_{\mathrm{R}} + N_{\mathrm{R}}^{\mathrm{c}} \end{pmatrix}, \\ M_{1,2} &= \frac{1}{2}(\sqrt{M_{\mathrm{R}}^2 + 4m_{\mathrm{D}}^2} \mp M_{\mathrm{R}}), \\ \tan\theta &= \frac{M_1}{m_{\mathrm{D}}} = (\frac{M_2}{m_{\mathrm{D}}})^{-1} \qquad (M_1 < M_2), \end{aligned}$$

where $s = \sin\theta, c = \cos\theta$. In the fourth generation the smaller mass eigenvalue M_1 should be greater than 45 GeV. Moreover, the Dirac mass $m_{\rm D}$ is bounded above because of the requirement of triviality; $m_{\rm D}$ is of the order of the weak scale. Consequently, it turns out that the Majorana mass $M_{\rm R}$ cannot be so large, and the two mass eigenvalues M_1 and M_2 are also the same of order of magnitude, *i.e.*, the see-saw is almost balancing in the heavy neutrino sector. In this case, the existence of the right-handed Majorana mass $M_{\rm R}$ affects various weak processes through the large left-right mixing. Contributions to S and T parameters in this model have already been calculated and turned out to be negative when $\tan\theta \sim 0.4$ [7, 8]. We will pay attention to this Majorana mass region, because the precision mesurements at LEP favor negative values for S and T [9].

With the self-conjugacy of the majorana fields N_1 and N_2 under the charge conjugation operation, the fermion currents can be written in terms of mass-eigenstates,

$$\begin{aligned}
J^Q_\mu &= Q^{\rm E}\overline{E}\gamma_\mu E, \\
J^3_\mu &= I^{\rm E}_3\overline{E}\gamma_\mu\frac{1-\gamma_5}{2}E + I^{\rm N}_3\left(-\frac{1}{2}c^2\overline{N_1}\gamma_\mu\gamma_5 N_1 - \frac{1}{2}s^2\overline{N_2}\gamma_\mu\gamma_5 N_2 + isc\overline{N_2}\gamma_\mu N_1\right), \\
J^+_\mu &= \frac{1}{\sqrt{2}}\left(-ic\overline{N_1} + s\overline{N_2}\right)\gamma_\mu\frac{1-\gamma_5}{2}E, \\
J^-_\mu &= \frac{1}{\sqrt{2}}\overline{E}\gamma_\mu\frac{1-\gamma_5}{2}\left(icN_1 + sN_2\right), \qquad (1)
\end{aligned}$$

where E refers to the charged lepton field. We note that the Z coupling to neutrinos with the same masses is axial vector, while it becomes a vector coupling when mixing the neutrinos with different masses.

We now consider the radiative corrections to $\mathrm{e^+e^-} \longrightarrow \mathrm{W^+W^-}$ at one-loop level. Because the mixing between heavy leptons and an electron is probably small and negligible, if it exist, the radiative corrections appear dominantly in the self energies and the corrections to the trilinear vertex corrections of gauge bosons. These corrections belong to “oblique” ones [1]. A renormalization program for oblique corrections is proposed by Kennedy and Lynn on the concept of effective lagrangian [1]. The extension of their formalism to the process of $\mathrm{e^+e^-} \longrightarrow \mathrm{W^+W^-}$ is given by Ahn, Peskin, Lynn and Selipsky [4]. They express the s-channel amplitude at one-loop level for on-shell W bosons with running coupling constant e_*, the running wave-function renormalization constant of W boson $Z_{\rm W*}$ [1] and the vertex function $\Gamma^{\mu\alpha\beta}$ into which A and Z vertices are combined:

$$\mathcal{M}^{\rm 1-loop}_s = -\frac{ie^2_*}{P^2}Z_{\rm W*}\overline{v}\gamma_\mu u\Gamma^{\mu\alpha\beta}\varepsilon^*_\alpha(q)\varepsilon^*_\beta(\overline{q}),$$

where P, q and $\overline{q}$ are momenta of V, $\mathrm{W^-}$ and $\mathrm{W^+}$, u and v are spinors of $\mathrm{e^-}$ and $\mathrm{e^+}$ and $\varepsilon_\alpha(q)$ and $\varepsilon_\beta(\overline{q})$ refer to the polarization-vectors of $\mathrm{W^-}$ and $\mathrm{W^+}$, respectively.

Here, we define the modified vertex function $\overline{\Gamma}^{\mu\alpha\beta} = Z_{\rm W*}\Gamma^{\mu\alpha\beta}$ and express it in terms of W-pair form factors $f_i^{\rm V=A,Z}$ and canonical Lorentz structures T_i, following Hagiwara *et al.* [10]. (In ref. [4], $Z_{\rm W*}$ is omitted in the definition of W-pair form factors and is multiplied to the cross sections afterward as the overall factor.)

$$\overline{\Gamma}^{\mu\alpha\beta} = \sum_{i=1}^{7}(Qf_i^{\rm A} + \frac{I_3 - s_*^2 Q}{s_*^2}\frac{P^2}{P^2 - m_{\rm Z}^2}f_i^{\rm Z})T_i^{\mu\alpha\beta},$$

$$\begin{aligned}
T_1^{\mu\alpha\beta} &= (q-\overline{q})^\mu g^{\alpha\beta}, & T_2^{\mu\alpha\beta} &= -\frac{1}{m_{\rm W}^2}(q-\overline{q})^\mu P^\alpha P^\beta, \\
T_3^{\mu\alpha\beta} &= P^\alpha g^{\mu\beta} - P^\beta g^{\mu\alpha}, & T_4^{\mu\alpha\beta} &= i(P^\alpha g^{\mu\beta} + P^\beta g^{\mu\alpha}), \\
T_5^{\mu\alpha\beta} &= i\epsilon^{\mu\alpha\beta\rho}(q-\overline{q})_\rho, & T_6^{\mu\alpha\beta} &= -\epsilon^{\mu\alpha\beta\rho}P_\rho, \\
T_7^{\mu\alpha\beta} &= -\frac{1}{m_{\rm W}}(q-\overline{q})^\mu\epsilon^{\alpha\beta\rho\sigma}P_\rho(q-\overline{q})_\sigma,
\end{aligned}$$

where s_* is the running weak mixing angle defined by Kennedy and Lynn, Q and I_3 are the charge and the isospin of e$^-$, respectively, and we use the Minkowskian convention with $\epsilon_{0123} = 1$. Form factors can be written in terms of the self energies Π_{XY} and the vertex corrections Σ_X^i

$$\begin{aligned}
f_1^{\rm A} &= 1 + g_*^2[\Sigma_Q^1 + \Pi'_{11}], \\
f_1^{\rm Z} &= 1 + \frac{g_*^2}{c_*^2}[(\Sigma_3^1 - s_*^2\Sigma_Q^1) + c_*^2\Pi'_{11}] + \frac{M_{\rm Z*}^2 - m_{\rm Z}^2}{P^2 - m_{\rm z}^2}, \\
f_3^{\rm A} &= 2 + g_*^2[\Sigma_Q^3 + 2\Pi'_{11}], \\
f_3^{\rm Z} &= 2 + \frac{g_*^2}{c_*^2}[(\Sigma_3^3 - s_*^2\Sigma_Q^3) + 2c_*^2\Pi'_{11}] + 2\frac{M_{\rm Z*}^2 - m_{\rm Z}^2}{P^2 - m_{\rm z}^2}, \\
f_i^{\rm A} &= g_*^2[\Sigma_Q^i], \\
f_i^{\rm Z} &= \frac{g_*^2}{c_*^2}[(\Sigma_3^i - s_*^2\Sigma_Q^i)], \qquad i = 2,\, 4,\, 6,\, 7,
\end{aligned} \tag{2}$$

where we adopt the notation of ref. [4], g_* and $M_{\rm Z*}$ is the running coupling constant and the running Z boson's mass defined by Kennedy and Lynn, respectively, and $\Pi'_{11} \equiv \Pi'_{11}(m_{\rm W}^2)$. These form factors, especially in the limit of low energies, are controled by gauge or global symmetries [4, 10, 11, 12].

Using the currents in eq.(1), we can now calculate the one-loop corrections in eq.(2) due to the leptons in the fourth generation. First, we present the expressions for the self energies

$$16\pi^2\Pi_{QQ} = -(Q^{\rm E})^2 8P^2[\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_{\rm E}^2, M_{\rm E}^2)],$$

$$
\begin{aligned}
16\pi^2\Pi_{3Q} &= -(Q^{\mathrm{E}}I_3^{\mathrm{E}})4P^2[\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2)], \\
16\pi^2\Pi_{33} &= -(I_3^{\mathrm{E}})^2[4P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2)\} - 2M_{\mathrm{E}}^2\{\overline{\epsilon}^{-1} + \mathbf{b_0}(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2)\}] \\
&\quad - (I_3^{\mathrm{N}}c^2)^2[4P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_1^2, M_1^2)\} - 4M_1^2\{\overline{\epsilon}^{-1} + \mathbf{b_0}(P^2, M_1^2, M_1^2)\}] \\
&\quad - (I_3^{\mathrm{N}}s^2)^2[4P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_2^2, M_2^2)\} - 4M_2^2\{\overline{\epsilon}^{-1} + \mathbf{b_0}(P^2, M_2^2, M_2^2)\}] \\
&\quad - 2(I_3^{\mathrm{N}}c^2)(I_3^{\mathrm{N}}s^2)[4P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_1^2, M_2^2)\} \\
&\qquad\qquad -(M_1 - M_2)^2\{\overline{\epsilon}^{-1} + \mathbf{b_0}(P^2, M_1^2, M_2^2)\} \\
&\qquad\qquad +(M_1^2 - M_2^2)\{\mathbf{b_1}(P^2, M_1^2, M_2^2) - \mathbf{b_1}(P^2, M_2^2, M_1^2)\}], \\
16\pi^2\Pi_{11} &= -c^2[2P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_1^2, M_{\mathrm{E}}^2)\} - \frac{1}{2}(M_{\mathrm{E}}^2 + M_1^2)\overline{\epsilon}^{-1} \\
&\qquad\qquad +M_1^2\mathbf{b_1}(P^2, M_1^2, M_{\mathrm{E}}^2) + M_{\mathrm{E}}^2\mathbf{b_1}(P^2, M_{\mathrm{E}}^2, M_1^2)\}] \\
&\quad - s^2[2P^2\{\frac{1}{6}\overline{\epsilon}^{-1} - \mathbf{b_3}(P^2, M_2^2, M_{\mathrm{E}}^2)\} - \frac{1}{2}(M_{\mathrm{E}}^2 + M_2^2)\overline{\epsilon}^{-1} \\
&\qquad\qquad +M_2^2\mathbf{b_1}(P^2, M_2^2, M_{\mathrm{E}}^2) + M_{\mathrm{E}}^2\mathbf{b_1}(P^2, M_{\mathrm{E}}^2, M_2^2)\}],
\end{aligned}
\tag{3}
$$

where $\overline{\epsilon}^{-1}$ is the divergence of the dimentional regulatization, $\overline{\epsilon}^{-1} = 2/(4-D) - \gamma + \ln 2\pi$. The vertex corrections can be written as follows

$$
\begin{aligned}
\Sigma_Q &= \frac{1}{2}Q^{\mathrm{E}}c^2[H(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_1^2) - G(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_1^2)] \\
&\quad + \frac{1}{2}Q^{\mathrm{E}}s^2[H(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_2^2) - G(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_2^2)], \\
\Sigma_3 &= \frac{1}{2}I_3^{\mathrm{E}}[c^2H(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_1^2) + s^2H(P^2, M_{\mathrm{E}}^2, M_{\mathrm{E}}^2, M_2^2)] \\
&\quad - \frac{1}{2}I_3^N[c^4\{\tilde{H}(P^2, M_1^2, M_1^2, M_{\mathrm{E}}^2) + \tilde{G}(P^2, M_1^2, M_1^2, M_{\mathrm{E}}^2)\} \\
&\qquad\qquad +s^4\{\tilde{H}(P^2, M_2^2, M_2^2, M_{\mathrm{E}}^2) + \tilde{G}(P^2, M_2^2, M_2^2, M_{\mathrm{E}}^2)\} \\
&\qquad\qquad +2s^2c^2\{\tilde{H}(P^2, M_1^2, M_2^2, M_{\mathrm{E}}^2) - \tilde{G}(P^2, M_1^2, M_2^2, M_{\mathrm{E}}^2)\}].
\end{aligned}
\tag{4}
$$

The functions H and G in the above expression are

$$
\begin{aligned}
16\pi^2H(P^2, m_1^2, m_2^2, m_3^2) &= T_0[-\frac{2}{3}\overline{\epsilon}^{-1} + \mathbf{c_0} + \mathbf{c_1} - \frac{m_{\mathrm{W}}^2}{\mu^2}(\mathbf{c_4} - \mathbf{c_5}) - \frac{P^2}{\mu^2}(\mathbf{c_6} + \mathbf{c_7})] \\
&\quad + T_2[-4\frac{m_{\mathrm{W}}^2}{\mu^2}\mathbf{c_7}] \\
&\quad + T_3[\mathbf{c_0} - 3\mathbf{c_1} - \frac{m_{\mathrm{W}}^2}{\mu^2}(2\mathbf{c_3} - 5\mathbf{c_4} + 3\mathbf{c_5}) + \frac{P^2}{\mu^2}(\mathbf{c_6} + 3\mathbf{c_7})] \\
&\quad + T_5[\mathbf{c_0} - 3\mathbf{c_1} + \frac{m_{\mathrm{W}}^2}{\mu^2}(\mathbf{c_4} - \mathbf{c_5}) - \frac{P^2}{\mu^2}(\mathbf{c_6} - \mathbf{c_7})],
\end{aligned}
$$

$$16\pi^2 G(P^2, m_1^2, m_2^2, m_3^2) = (T_0 - T_3 + T_5)[\frac{m_1 m_2}{\mu^2}(\mathbf{c_2} - \mathbf{c_3})]. \tag{5}$$

Here $T_0 \equiv T_1 + 2T_3$ is the only Lorentz structure at tree level, μ is a reference point and the $\mathbf{c_i}$ have arguments $(P^2, m_1^2, m_2^2, m_3^3)$. We ommit gauge anomaly terms in eq.(5), because they should cancel themselves if we consider the full fourth generation. The functions $\tilde{H}$ and $\tilde{G}$ are obtained by reversing the sign of T_5 in eq.(5). The reduced Passarino-Veltman functions $\mathbf{b_i}(P^2, m_1^2, m_2^2)$ and $\mathbf{c_i}(P^2, m_1^2, m_2^2, m_3^3)$ in eq.(3) and (5) [4, 13] are defined as folows

$$\begin{aligned}
[\mathbf{b_0}, \mathbf{b_1}, \mathbf{b_3}] &= \int dxdy\delta(1-x-y)\ln(D_2/\mu^2)[-1, x, xy], \\
[\mathbf{c_0}, \mathbf{c_1}] &= \int dxdydz\delta(1-x-y-z)\ln(D_3/\mu^2)[1, z], \\
[\mathbf{c_2}, \mathbf{c_3}, \mathbf{c_4}, \mathbf{c_5}, \mathbf{c_6}, \mathbf{c_7}] &= \int dxdydz\delta(1-x-y-z)(\mu^2/D_3) \\
&\quad \times[1, z, z^2, z^3, xy, xyz],
\end{aligned}$$

$$\begin{aligned}
D_2 &= -xyP^2 + xm_1^2 + ym_2^2 - i\epsilon, \\
D_3 &= -z(1-z)m_{\mathrm{W}}^2 - xyP^2 + xm_1^2 + ym_2^2 + zm_3^2 - i\epsilon.
\end{aligned} \tag{6}$$

We note that Lorentz structures T_4, T_6 and T_7 disappear in eq.(5). The W-pair form factors corresponding to these Lorentz structures are CP variant [10], and hence should vanish explicitly in CP conserving models such as we consider here. We also note that the threshold behaviour of the form factors for N_1s' (N_2's) pair production and for N_1-N_2 production are different. In the self energies, this difference can be seen easily from eq.(3), as a differnce in the power of β_{ij}; $\beta_{ij} \equiv [1 - (M_i + M_j)^2/P^2]^{1/2}$. The leading terms of Π_{33} at N_1s' (N_2's) pair production threshold and at N_1-N_2 production threshold are proportional to $\beta^3_{11(22)}$ and β_{12}, respectively [14]. We will discuss the threshold behaviour of the form factors containing the vertex corrections afterward.

Now, we peform the numerical calculations using the method in ref [15]. We fix the following parameters to the values at energy scale of m_{Z}:

$$\alpha_*^{-1} = \frac{4\pi}{e_*^2} = 128.0, \qquad s_*^2 = 0.2319, \qquad m_{\mathrm{Z}} = 91.187 \text{ [GeV]},$$

where we neglect the running effects of e_*, s_* [1, 2]. We also set $m_{\mathrm{W}} = 80.22$ [GeV].

The real parts of the corrections to W-pair form factors $\Delta f^{\mathrm{Z}}_{i=1,2,3,5}$ from heavy neutrinos and charged leptons are shown in fig.1(a) and 1(b), where neutrinos are Dirac and Majorana particles, respectively. The form factors $\Delta f_2^{\mathrm{A,Z}}$ are quite small, as can be seen in fig.1, since these form factors correspond to the dimension 6 operators [4, 10]. Therefore the radiative corrections due to the heavy fermions mainly appear in the other form factors. We can see cusps in the $\Delta f^{\mathrm{Z}}_{1,3,5}$-lines at the Dirac neutrino threshold in fig.1(a) and in the $\Delta f^{\mathrm{Z}}_{5}$-line at N_1-N_2 threshold in fig.1(b). The enhancement at N_1-N_2 threshold is very small because of the small couplings including the mixing angle in eq.(1). On the other hand, $\Delta f^{\mathrm{Z}}_{1,3,5}$-lines are not cusped at N_1-N_1 threshold as seen in fig.1(b). In self energies, as mentioned above, the difference of threshold behaviours depending on the neutrino types has been checked analytically. Numerical calculations including vertex corrections also suggest that threshold behaviours are scalar-like for N_1's (N_2's) pair production, but Dirac fermion-like for N_1-N_2 production. This is caused by the difference in the coupling types between $Z\overline{N_1}N_1$ ($Z\overline{N_2}N_2$) and $Z\overline{N_1}N_2$ in eq.(1). It is also helpful to recognize that the two Majorana particles of the same kind (N_1-N_1 or N_2-N_2) are produced by the p-wave from the spin 1 photon or Z boson, in the same manner as the scalars' pair production, whereas the different kinds of neutrinos (N_1-N_2) are produced by the s-wave, which happens in the ordinary Dirac fermions' pair production. The difference appeared in the threshold behaviours may be understood as β^{2l+1} for the l-th wave with the relative velocity β.

Next, we present the results on the differential cross sections $d\sigma/d\cos\Theta$ and the forward-backward asymmetries A_{FB}. Since the vertex corrections, on which we concentrate in this paper, appear only on s-channel gauge boson exchange diagrams, we plot $d\sigma/d\cos\Theta$ and A_{FB} for right-handed polarized electrons in fig.2 $\sim$ fig.4. Hereafter we consider the full fourth generation with degenerate quarks, and set $M_{\mathrm{U}} = M_{\mathrm{D}} = M_{\mathrm{E}} = 500$[GeV], where M_{U}, M_{D} and M_{E} are masses of up-type quark, down-type quark and charged lepton, respectively. In fig.2, we plot the differential cross section $d\sigma/d\cos\Theta$ at scattering angle $\Theta = \frac{\pi}{2}$ in units of the point cross section $1R = 4\pi\alpha_*^2/3s$ with $s = P^2$ and $\alpha_* = 1/128.0$. We find that the differntial cross sections depend on the neutrino types, i.e., Majorana neutrino or Dirac neutrino, and their masses, although no sizable peaks can be seen except for the charged particles' pair production threshold. The anguler distribution of the differntial cross section is shown in fig.3. At any scattering angle, the heavy fermions make the correction less than about 1% at $\sqrt{s} = 170$ [GeV] and less than about 4% at $\sqrt{s} = 300$ [GeV]. In fig.4 we plot the forward-backward asymmetry A_{FB} for $\mathrm{e}^+_{\mathrm{L}}\mathrm{e}^-_{\mathrm{R}} \to \mathrm{W}^+\mathrm{W}^-$ with unpolarized final W's, in the angular range $|\cos\Theta| \leq 0.98$. For this initial polarization , A_{FB} is controled by two form factors f^{A}_5 and f^{Z}_5 which vanish at tree level. At one-loop level, these form factors also vanish when we consider a full generation with degenerate Dirac fermions. (Note that the functions $\tilde{H}$ and $\tilde{G}$ in eq.(4) are obtained by reversing the sign of T_5 in eq.(5).) Hence the non-vanishing A_{FB} is solely due to the Majorana mass M_{R} in line(B) or due to the Dirac mass spliting in line(D).

To discuss the observability of the radiative corrections at LEP200 or at the next generation linear colliders, we estimate the cross section for unpolarized electrons. At tree level, the cross section σ in the angular range $|\cos\Theta| \leq 0.98$ is 16.80 [pb] at $\sqrt{s} = 175$ [GeV] and 11.45 [pb] at $\sqrt{s} = 300$ [GeV]. The effect of radiative corrections is $0.6 \sim 0.7\%$ at both energies. At LEP200 with an integrated luminosity of 500 [pb^{-1}], it is difficult to measure the deviation from the tree level value. However, at the next generation linear colliders with $\sqrt{s} = 300\,[\mathrm{GeV}] \sim 1.5\,[\mathrm{TeV}]$ and an integrated luminosity of 5×10^4 [pb^{-1}], the deviation depend on the neutrino types and their masses will be mesureable.

In summary, we have investigated the one-loop corrections due to the heavy Majorana neutrinos in the process $\mathrm{e}^+\mathrm{e}^- \to \mathrm{W}^+\mathrm{W}^-$ which will be seen in LEP200 and the next generation linear colliders. We have estimated W-pair form factors in terms of reduced Passarino-Veltman functions. We have found that the threshold behaviour of these form factors is scalar-like for a pair of light neutrinos (N_1-N_1) or a pair of heavy neutrinos (N_2-N_2) production, while that is Dirac fermion-like for light and heavy neutrinos (N_1-N_2) production. We also have estimated the differential cross section and the forward-backward asymmetry, numerically, and have found them depending on the Dirac mass and the mixing angle in the see-saw mechanism even at low energies. The deviation of the cross section from the tree level one will be measurable at the next linear colliders.

Acknowledgments

The authors would like to thank N. Nakazawa and K. Kato who offered the computer program useful for performing Feynman Parameter integrals. They are also grateful to T. Inami, C.S. Lim and I.Watanabe for discussions. This work is in part supported by Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (No. 05640336).

References

[1] D. C. Kennedy and B. W. Lynn, *Nucl. Phys.* **B322** (1989) 1

[2] M. E. Peskin and T. Takeuchi, *Phys. Rev. Lett.* **65** (1990) 964; *Phys. Rev.* **D46** (1992) 381

[3] G. Altarelli and R. Barbieri, *Phys. Lett.* **B253** (1990) 161;
G. Altarelli, R. Barbieri and S. Jadach *Nucl. Phys.* **B369** (1992) 3

[4] C. Ahn, M. E. Peskin, W. Lynn and S. Selipsky, *Nucl. Phys.* **B309** (1988) 221

[5] M. Gell-Mann, P. Ramond and R. Slansky, in : *Supergravity*,
eds. P. V. Nieuwenhuizen and D. Z. Freedman (North Holland, Amsterdam, 1979) p.315
T. Yanagida, Horizontal Gauge Symmetry and Masses of Neutrinos,
in: *Proc. Workshop on Unified theories and baryon number in the universe* (KEK, February 1979), eds. O. Sawada and A. Sugamoto (KEK-79-18) p.95
R. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44** (1980) 912; *Phys. Rev.* **D23** (1981) 165

[6] C. T. Hill and E. A. Paschos, *Phys. Lett.* **B241** (1990) 96

[7] S. Bertolini and A. Sirlin, *Phys. Lett.* **B257** (1991) 179

[8] E. Gates and J. Terning, *Phys. Rev. Lett.* **67** (1991) 1840

[9] D. C. Kennedy and P. Langacker, *Phys. Rev. Lett.* **65** (1990) 2967; **66** (1991) 395(E);
W. J. Marciano and J. Rosner, *Phys. Rev. Lett.* **65** (1990) 2963

[10] K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, *Nucl. Phys.* **B282** (1987) 253

[11] B. Holdom, *Pyhs. Lett.* **B258** (1991) 156

[12] T. Appelquist and G.H. Wu, *Phys. Rev.* **D48** (1993) 3235

[13] G. Passarino and M. Veltman, *Nucl. Phys.* **B160** (1979) 151

[14] M. Marui, preprint OCHA-PP 35 (1993)

[15] J. Fujimoto, Y. Shimizu, K. Kato and Y. Oyanagi, *Progr. Theor. Phys.* **87** (1992) 1233

Figure Captions

fig.1 Correction $\Delta f^{\mathrm{Z}}_{i=1,2,3,5}$ to W-pair form factors due to the fourth generation leptons as a function of $\sqrt{s}$.

Fig.1(a) is given for the Dirac neutrino N with $M_{\mathrm{N}} = 200$ [GeV]. Fig.1(b) is given for the Majorana neutrinos N_1 and N_2 with $M_1 = 200$ [GeV] and $M_2 = 1250$ [GeV] ($m_{\mathrm{D}} = 500$ [GeV] and $\tan\theta = 0.4$). We set the charged lepton mass M_{E} at 500 [GeV] in the both figure.

fig.2 Differntial cross section $\frac{d\sigma}{d\cos\Theta}$ for $\mathrm{e}^+_{\mathrm{L}}\mathrm{e}^-_{\mathrm{R}} \to \mathrm{W}^+\mathrm{W}^-$ at scattering angle $\Theta = \frac{\pi}{2}$ as a function of $\sqrt{s}$.

The dotted line (A) shows the differntial cross section at tree level. The solid line (B), the dashed-and-dotted line (C) and the dashed line (D) include the corrections due to the quarks and the leptons of the fourth generation at one-loop level. The line (B) is depicted for the Majorana neutrinos N_1 and N_2 with $M_1 = 200$ [GeV] and $M_2 = 1250$ [GeV]. The line (C) and the line (D) are shown for the Dirac neutrino N with $M_{\mathrm{N}} = 500$ [GeV] and 200 [GeV], respectively.

fig.3 Angular distribution of the differntial cross section $d\sigma/d\cos\Theta$ for $\mathrm{e}^+_{\mathrm{L}}\mathrm{e}^-_{\mathrm{R}} \to \mathrm{W}^+\mathrm{W}^-$. Fig.3(a) and 3(b) are at $\sqrt{s} = 175$ [GeV] and 300 [GeV], respectively. The values of the masses in (A),(B),(C) and (D) are the same as those in the corresponding curves in fig.2.

fig.4 Forward-backward asymmetry

$$A_{FB} = \frac{\int_0^{0.98} d\cos\Theta \frac{d\sigma}{d\cos\Theta} - \int_{-0.98}^{0} d\cos\Theta \frac{d\sigma}{d\cos\Theta}}{\int_0^{0.98} d\cos\Theta \frac{d\sigma}{d\cos\Theta} + \int_{-0.98}^{0} d\cos\Theta \frac{d\sigma}{d\cos\Theta}}$$

for $\mathrm{e}^+_{\mathrm{L}}\mathrm{e}^-_{\mathrm{R}} \to \mathrm{W}^+\mathrm{W}^-$ as a function of $\sqrt{s}$.

The values of the masses in (A), (B), (C) and (D) are the same as those in the correponding curves in fig.2.



M_E =500 [GeV]
M_N =200 [GeV]
: Δf_1^Z
: Δf_2^Z
: Δf_3^Z
: Δf_5^Z
0.004
0.002
0.000
−0.002
Δf_i^Z
500
1000
1500
2000
$\sqrt{s}$ [GeV]

Fig.1(a)



M_E =500 [GeV]
M_1 =200 [GeV]
M_2 =1250[GeV]
: Δf_1^Z
: Δf_2^Z
: Δf_3^Z
: Δf_5^Z
0.004
0.002
0.000
−0.002
Δf_i^Z
500
1000
1500
2000
$\sqrt{s}$ [GeV]

Fig.1(b)


0.45
0.40
0.35
0.30
0.25
0.20
0.15
0
500
1000
1500
2000
$\sqrt{s}$ [GeV]
$d\sigma/d\cos\Theta$ [R]
: (A)
: (B)
: (C)
: (D)

Fig.2


0.0775
0.0750
0.0725
0.0700
0.0675
0.0650
dσ/dcosΘ [R]
−1
−0.5
0
0.5
1
cosΘ
√s =175[GeV]
: (A)
: (B)
: (C)
: (D)

Fig.3(a)



Fig.3(b)



Fig.4